

# Adverse Childhood Experiences in Jefferson County, Indiana

2021 IU Southeast Sociology Research Lab

## ACE: ADVERSE CHILDHOOD EXPERIENCES

Before age 18

LIVED WITH ANYONE WHO WAS...

a problem drinker or alcoholic or who used street drugs




LIVED WITH ANYONE WHO WAS...

depressed, mentally ill, or suicidal



LIVED WITH ANYONE WHO...

was sentenced to serve time in a prison, jail or other correctional facility



HAD NO ONE IN YOUR FAMILY WHO

loved you or thought you were important or special



HAD PARENTS WHO WERE...

separated or divorced



HAD A MOTHER OR STEPMOTHER WHO WAS OFTEN...

slapped, hit, grabbed, kicked, pushed or had things thrown at her



HAD PARENTS OR ADULTS IN YOUR HOME WHO OFTEN...

hit, beat, kicked, or physically hurt you in any way before you were age 18



HAD A PARENT OR ADULT IN YOUR HOME WHO OFTEN...

swore at you, insulted you, or put you down



HAD ANYONE AT LEAST 5 YEARS OLDER THAN YOU OR AN ADULT WHO OFTEN...

touched you sexually or tried to make you touch sexually



HAD ANYONE AT LEAST 5 YEARS OLDER THAN YOU OR AN ADULT WHO OFTEN...

forced you to have sex



<https://www.resilientjeffersoncounty.org/>

Adverse Childhood Experiences (ACEs) are traumatic events experienced from birth through age 17. Research shows that these experiences disrupt neurodevelopment, causing social, emotional, and cognitive impairments that affect behaviors, including health behaviors.<sup>1</sup> These impacts lead to negative outcomes in health, mental health, education, and social success and well being, ultimately resulting in early death.

Jefferson County is home to relatively high rates of smoking, obesity, physical inactivity, poor mental and physical health days, and diabetes. In addition, the county has a far higher suicide rate than the state or nation. Jefferson County has relatively low educational attainment, and the median household income is below the state median. In the face of these challenges, identifying leverage points for improving outcomes is vitally important.


The Jefferson County community-wide ACEs survey is an attempt to document the prevalence of ACEs, draw connections between experiences of childhood trauma and health, mental health, education, and economic outcomes, and identify the highest priorities for prevention and intervention.

## Key Findings

Based on a weighted sample:

- The estimated average ACE score for Jefferson County adults is 2.4.
- 30.4% of Jefferson County adults have an ACE score of 4 or higher.
- Roughly 9.7% of Jefferson County adults have an ACE score of 7 or higher.
- More than a third of Jefferson County adults report that before age 18 they experienced one or more of the following: separation or divorce of parents, emotional abuse, or living with someone who suffered from substance use disorder (an alcoholic or drug user).
- 32.2% of Jefferson County adults have no ACEs.

Figure 1: Impact of ACEs Throughout the Lifespan



Source: CDC-Kaiser Health. 2020[1998]. "About the CDC-Kaiser ACE Study." *Violence Prevention* (<https://www.cdc.gov/violenceprevention/ACE/about.html>).

## Research on ACEs

In 1998, Vincent Felitti, Robert Anda, and a team of researchers published a groundbreaking study documenting statistically significant relationships between experiences of seven categories of childhood trauma and health risk behaviors and disease in adulthood.<sup>2</sup> Since that time, dozens of studies have replicated and expanded on the initial findings, confirming consistent relationships between ACEs and negative outcomes in the areas of health, mental health, substance abuse, educational attainment, employment stability, and income. This field of research finds strong dosage effects where those with higher ACE scores experience progressively higher rates of disease and dysfunction in their adult lives. Those with zero reported ACEs have significantly better outcomes than those with four or more ACEs across a range of indicators of adult well-being and predictors of early death.<sup>3</sup>

## Sex, Age, Education, & Income

The inclusion of ACEs as part of the Behavioral Risk Factor Surveillance System (BRFSS) has allowed researchers to amass a very large geographically diverse sample of adults from 34 states. While childhood trauma is common across sociodemographic groups, the research suggests that some populations are at risk for higher ACE scores than others.<sup>4</sup>

A majority of American adults (57.8%) experienced at least one childhood trauma and 21.5% experienced 3 or more ACEs.<sup>5</sup> Multiracial individuals have a significantly higher average ACE score (2.4) than all other races and ethnicities, and whites have the lowest average ACE score of the racial categories (1.5).<sup>6</sup> Women have a higher mean ACE score than men (1.6 compared to 1.5), and those age 25-34 report a significantly higher mean ACE score than any other group (2.0).<sup>7</sup>

Those with higher income and educational attainment have lower ACE scores than those with lower income and educational attainment.<sup>8</sup>

Research suggests that among the mechanisms by which ACEs produce poorer outcomes in education and employment are the health and mental health outcomes that affect work performance and stability.<sup>9</sup>

Research in Wales suggests the impact of ACEs on educational attainment is largely explained by failure to complete high school.<sup>10</sup> Research indicates that a combination of lower educational attainment and challenges with employment stability explain negative income outcomes for those with higher ACE scores.<sup>11</sup>

## Health

Toxic stress is one mechanism by which childhood trauma affects adult health and mental health outcomes. When the body experiences chronic stressors, it remains in a fight or flight mode of vigilance that causes the release of hormones and chemicals that affect neurodevelopment and long-term health outcomes.<sup>12</sup> Chronic stress causes changes to neurobiology that impact early brain development,<sup>13</sup> the immune system,<sup>14</sup> and the endocrine system.<sup>15</sup> Stressful or traumatic experiences often lead to social, emotional, and cognitive deficiencies that increase risk for unhealthy behaviors and chronic disease.<sup>16</sup>

Toxic stress can cause issues with immune and metabolic systems that fight illness, leading to a

---

lifetime of susceptibility to illness.<sup>17</sup> Toxic stress causes early onset of disease, disability, and premature death.<sup>18</sup> ACE exposure increases risk of obesity.<sup>19</sup> Higher ACE scores are associated with increased risk of smoking, heart disease, chronic lung disease, poor self-rated health, having 50 or more sexual intercourse partners, and sexually transmitted disease (2-4 fold increase in risks).<sup>20</sup> Researchers find a 1.4-1.6 fold increase in physical inactivity and severe obesity among those with 4 or more ACEs compared to those with none.<sup>21</sup> The number of ACEs showed a dose dependent relationship with diseases such as ischemic heart disease, cancer, chronic lung disease, skeletal fractures, and liver disease.<sup>22</sup> Research links ACEs to systemic arterial stiffness among adolescents.<sup>23</sup>

## Mental Health and Substance Use Disorder

Childhood trauma increases risk for a range of mental health concerns and for substance use disorder (SUD). Bryant (2020) found that each individual ACE type was significantly associated with SUD.<sup>24</sup> These issues may impact employment stability. Studies find associations between early adversity and each of the following:

- Frequent mental distress.<sup>25</sup>
- Depression.<sup>26</sup>
- Suicide attempts.<sup>27</sup>
- Smoking.<sup>28</sup>
- Alcohol abuse.<sup>29</sup>
- Substance abuse.<sup>30</sup>

In addition, sexual abuse and parental/other family member's mental illness increase the odds for having a suicide attempt for both men and women and emotional neglect is also a factor for men.<sup>31</sup>

While some may understand ACEs as individual level experiences that shape individual outcomes, social connectedness, community, and public policy may play significant roles in preventing and mitigating the impacts of adverse childhood experiences.<sup>32</sup> Moreover, a high prevalence of ACEs results in community level impacts such as lower educational attainment, labor force issues, poor performance in indicators of health and quality of life.

## Methods

The Indiana University Southeast Sociology Research Lab and Applied Research and Education Center (AREC) used the Centers for Disease Control's 10-item ACE instrument as the base for the Jefferson County community-wide ACEs survey. Based on the review of the literature, the team asked additional questions to obtain information on adult outcomes found to be associated with ACEs. In addition, the team included demographic questions to allow for comparisons between Jefferson County and national patterns.


The Indiana University Institutional Review Board reviewed and approved the study design and materials. The survey was made available electronically using Qualtrics survey software. Participants were recruited using email distribution of the link through local nonprofit organizations and other mailing lists using a snowball sampling approach, where recipients were free to share and distribute as they chose. In addition, social media posts encouraged participation and linked to the online survey.

Flyers were posted throughout the county and quarter sheet flyers were distributed via local organizations with a QR code that linked directly to the survey. The QR code also appeared on the survey collection boxes to make the online survey available in all locations where paper surveys were distributed. Paper surveys were made available at 10 sites throughout the community, including both rural and in-town locations, and venues serving low-income and otherwise diverse populations that tend to be underrepresented in online surveys. Spanish language paper surveys were made available in locations serving the Hispanic population and the online survey was available in both English and Spanish. Locked survey boxes were available for survey collection along with business reply envelopes that allowed respondents to mail completed surveys directly to the AREC.

Three surveys were received via direct mail and 46 paper surveys came in through drop boxes. Six hundred and one electronic surveys were completed for a total of 666 surveys. Of those, 642 answered all 10 questions about childhood trauma.

The research team used SPSS 28 to manage and analyze the data. In order to better estimate

Figure 2: Males and Females in Sample and Jefferson County Weighted Sample




Note: Weighted sample reflects county demographics for age, sex, education, and income.

population figures, the sample was weighted to county population figures for sex, age, income, and education. Cross tabs with appropriate significance tests were used to test group differences, along with t-tests for differences in mean ACE scores across groups.

This report presents basic descriptive analysis for both the actual and weighted samples to ensure transparency around the estimation procedures and to provide a clearer sense of the likely prevalence of ACEs and associated adult outcomes in Jefferson County, Indiana.

Figure 3: Age



## Sample and County Demographics

The convenience sampling strategy produced a large sample, but one that is not representative of the community's demographics. The sample is heavily skewed toward women (Figures 2), under-represents 18-24 year-olds and those over age 65 (Figure 3), over-represents

those with higher education (Figure 4), and those from higher income categories (Figure 5).

Jefferson County is 94.5% White, 2.1% Black, 0.9% Asian, 1.2% two or more races, and 2.8% Hispanic. The sample was 98.9% White and only 1.9% Black and 2.0% Hispanic. The small number of minority responses make weighting for race and Hispanic problematic.

The weighted sample uses responses received and demographics for the county to generate a data set that approximates the full population and its sex, age, education, and income distributions. For example, our sample was 22.0% male (Figure 2); weights for sex generate a sample that matches the

Figure 4: Educational Attainment





Figure 5: Income Distribution



county sex composition, which is 46.9% male (Figure 2). Because demographics are associated with ACEs, weighting produces a more accurate estimate of the prevalence of ACEs in the County.

## Findings

The estimated average ACE score for adults in Jefferson County is 2.4 (SD=2.5) compared to a

national mean of 1.6. In total, approximately 32.2% of Jefferson County adults have an ACE score of 0 and 67.7% of respondents have one or more ACEs compared to 57.8% nationally (Figure 7). An estimated 30.3% of Jefferson County adults report four or more ACEs and 9.6% report seven or more ACEs. Women have a significantly higher mean ACE score than men (2.7 compared to 2.1 with a median of 2 for women and 1 for men).

The highest prevalence ACEs are parents divorced or separated (43.1%), emotional abuse & fear for safety (37.0%), living with an alcoholic or drug user (33.6%), and household member mentally ill or attempted suicide (31.8%). The lowest prevalence ACE is household member that ever went to prison at 10.1% compared to 8.1% nationally (Figure 8).


In the Jefferson County weighted sample, educational attainment is significantly associated with ACE scores. Those with four or more ACEs are significantly less likely to have a college degree or higher, are less likely to have a high

Figure 6: Race and Hispanic Ethnicity

	Sample (N=645)	Jefferson County
<b>White alone</b>	97.5%	95.0%
<b>American Indian &amp; Alaska Native alone</b>	0.2%	0.4%
<b>Black alone</b>	.9%	2.2%
<b>Asian alone</b>	0.0%	0.9%
<b>Two or More Races</b>	1.4%	1.6%
<b>Hispanic or Latino/a</b>	2.0%	3.0%

Note: The weighted sample over-represents Whites and Hispanics. Still, the small number of Black and Hispanic responses would make weighting for race and Hispanic origin problematic.

Figure 7: Distribution of ACE Scores in Jefferson County Actual and Weighted Samples



Note: As you read the charts, weighted figures are likely closer to true population rates than sample figures.

school diploma or equivalency, and are more likely to have less than high school (Figure 9). The Jefferson County survey asked respondents if they have ever struggled to maintain consistent employment. Among those who report 4 or more ACEs, 19.2% of the weighted sample reported that they had compared to 10.1% of the weighted sample with zero ACEs. The association is statistically significant (Figure 10,  $p < .001$ ).

Struggles with employment stability for those with higher ACE scores are likely the result of challenges with health, mental health, and substance abuse. When combined with lower educational attainment, these issues yield lower income levels. In the Jefferson County weighted sample the pattern is notable in higher percentages of those with four or more ACEs at lower income levels and lower percentages at the highest income levels than for those with zero ACEs. The significantly lower income levels of those with higher ACE scores makes it more likely

that children in their households will also experience chronic stress associated with poverty.

### Health, Mental Health, and Substance Abuse

ACEs are associated with poor health behaviors and high risk practices such as smoking and substance use that increase the likelihood of chronic disease and premature death. County Health Rankings data indicate that Jefferson County has higher average numbers of days of poor mental and physical health than the state average. Residents report an average 5.2 poor mental health days per month compared to 4.7 for the state and 3.8 among U.S. top performers, and 4.2 poor physical health days per month compared to 4.0 for the state and 3.4 among U.S. top performers. The county has relatively high rates of smoking (24%), obesity (33%), and physical inactivity (25%).\*

(Continued on page 8)

\*County Health Rankings round data to whole numbers so we are unable to present figures to the first decimal place.

Figure 8: Prevalence of each ACE in Jefferson County Actual and Weighted Samples




Figure 9: Educational Attainment for those with Zero ACEs and those with 4 or More ACEs, weighted sample (p< .001)





Figure 10: Percent with Zero ACEs and Four or More ACEs that Report Having Struggled with Employment Stability (Sample N=637|Weighted N=24,391, p<.001)



The survey asked respondents to report on a selection of health behaviors, physical and mental health outcomes to explore their association with ACEs among Jefferson County residents. Those with four or more ACEs are significantly more likely than those with zero

ACEs to smoke or to have smoked regularly at some point in their life (Figure 12). In Jefferson County, a higher ACE score is significantly associated with higher rates of sexually transmitted infections, heart disease, chronic lung

*(Continued on page 10)*

Figure 11: Income Distribution for those with Zero ACEs and those With 4 or More ACEs, weighted sample (N=24,476, p<001)






Figure 12: Percent Indicating they Smoke or Have Ever Regularly Smoked Cigarettes by ACE score category (Sample N=642 | Weighted N=24,474, p<.001)






Figure 13: Prevalence of Health Issues for those with Zero ACEs and those with Four or More ACEs (Sample N=438 | Weighted N=25,567)



Statistically significant differences in prevalence between those with zero ACE and those with four or more ACE are indicated with asterisks. \*p<.05, \*\*p<.01, \*\*\*p<.001. Contrary to findings in the literature, associations with liver disease and obesity were not found to be statistically significant in the Jefferson County sample, with or without weighting.

Figure 14: Prevalence of Reported Mental Health Concerns by ACE Score, Sample (N=448)



disease, and physical disability that limits movement (Figure 13); all confirming findings in the literature on ACEs and adult health outcomes. Interestingly, the original ACEs research focused on obesity. In Jefferson County, however, obesity rates were not significantly different between those with zero and those with four or more ACEs.

The weighted sample indicates positive associations between higher ACE scores and mental health issues among Jefferson County adults as well. Figure 14 illustrates that as one moves up in ACE score category, so too does the likelihood of depression, anxiety, and suicidal thoughts. When those with four or more ACEs are compared to those with zero ACEs (Figure 15), prevalence of each mental health issue is significantly higher among those with 4 or more ACEs ( $p < .001$ ).

Finally, in a state and region beset with some of the worst of the opioid epidemic, those with four or more ACEs are 10 times more likely than those with zero ACEs both to report a diagnosis or

self-assessment of substance use disorder and to have ever misused prescription pain killers or to have used street opioids such as heroin (Figure 16).

Findings from Jefferson County confirm the existing research on the association between ACEs and adult health, mental health, and substance abuse outcomes. With a higher average ACE score than the nation, it is not surprising that Jefferson County also sees higher rates of chronic disease, mental health challenges, and substance use disorder.

## Discussion

Childhood trauma disrupts neurodevelopment, causing social, emotional, and cognitive impairments that affect behaviors, including health behaviors.<sup>33</sup> These impacts lead to negative outcomes in health, mental health, education, income, and social success and well being, ultimately resulting in early death.

Aggregated across a county, a higher than

*(Continued on page 12)*

Figure 15: Prevalence of Reported Mental Health Issues for those with Zero ACEs Compared to those with 4 or More ACEs





Figure 16: Percent Indicating any Substance Use Disorder and Misuse of Prescription Painkillers or Use of Street Opioids Such as Heroin among those with Zero ACEs and those with Four or More ACEs



---

average prevalence of such experiences results in myriad challenges from low educational attainment and employment instability to high rates of chronic disease, substance use disorder, mental health concerns and suicide.

While childhood trauma is not the only factor that shapes quality of life outcomes, it is one that we understand, is well-documented, and can be prevented and mitigated through building individual and community resilience. A strong focus on building healthy childhoods, social connections, and effective parenting skills can lay the foundation for improving outcomes. Communities need to engage business, government, and the nonprofit sector to align resources and influence to build support for access to mental health care, quality care and education for children birth to 18, and community building activities that reduce isolation, build social ties, and improve resilience.

Effective and sustainable community and economic development efforts can reduce household financial stressors that contribute to the prevalence of ACEs. Prevention and mitigation priorities may include the following<sup>34</sup>:

- Create and redesign policies and jobs that support workers with living wages.
- Build early care, school, and extracurricular youth programs that support families (birth to 18).
- Teach social-emotional skills with attention to safe and healthy relationships and healthy parenting and family dynamics.
- Connect youth to caring adults.
- Provide widespread access to physical and mental health care and family-centered treatment for substance use disorders.

The data from Jefferson County suggest significant opportunities to improve quality of life and community well-being through preventing and mitigating the impacts of childhood trauma. A focus on resilience and social support for healthy individuals and families can provide leverage for stronger and more sustainable community and economic development.

## References

(Full citations available at [www.resilientjeffersoncounty.org](http://www.resilientjeffersoncounty.org))

- <sup>1</sup> Felitti, Vincent J. et al. 1998; Anda, Robert F. et al. 2010. Verbitsky-Savitz et al. 2016.
- <sup>2</sup> Felitti, et al. 1998.
- <sup>3</sup> Giano, Wheeler, and Hubach. 2020.
- <sup>4</sup> Giano, Wheeler, and Hubach 2020.
- <sup>5</sup> Giano, Wheeler, and Hubach 2020.
- <sup>6</sup> Giano, Wheeler, and Hubach 2020.
- <sup>7</sup> Giano, Wheeler, and Hubach 2020.
- <sup>8</sup> Giano, Wheeler, and Hubach 2020.
- <sup>9</sup> DeVenter et al. 2020; Blodgett 2018; Hardcastle et al. 2018;
- <sup>10</sup> Hardcastle et al. 2018;
- <sup>11</sup> DeVenter et al. 2020; Metzler et al. 2017; Liu et al. 2012; Zielinski. 2009.
- <sup>12</sup> McRory et al. 2011; Gilbert et al. 2010. Shonkoff et al., 2009; Shonkoff & Garner 2012; Roth 2009; Szyf 2009
- <sup>13</sup> Ibid.
- <sup>14</sup> Boullier and Blair 2018; Danese 2009.
- <sup>15</sup> Ibid.
- <sup>16</sup> Hillis et al. 2021. Verbitsky-Savitz, et al. 2016.; Brown et al. 2009; Dube et al. 2003. Felitti et al. 1998.
- <sup>17</sup> Boullier and Blair 2018; McRory et al. 2011; Gilbert et al. 2010. Danese 2009; Shonkoff et al., 2009; Shonkoff & Garner 2012; Roth 2009; Szyf 2009; Bierhaus et al. 2003.
- <sup>18</sup> Fuller-Thomson 2016; Verbitsky-Savitz 2016; Gilbert 2015; Brown 2009; Chapman et al. 2004; Danese et al. 2009; Felitti et al. 1998
- <sup>19</sup> Felitti 1998; Verbitsky-Savitz 2016.
- <sup>20</sup> Felitti 1998; Verbitsky-Savitz 2016; Shonkoff et al. 2012; Wade et al 2012; Hillis et al 2001; Klassen et al. 2016
- <sup>21</sup> Anda et al. 2010. Minnesota Department of Health 2013; Felitti, et al. 1998.
- <sup>22</sup> Felitti 1998; Gilbert et al. 2015.
- <sup>23</sup> Klassen et al 2016.
- <sup>24</sup> Bryant, Coman, and Damian 2020.
- <sup>25</sup> Gilbert et al 2015.
- <sup>26</sup> Gilbert et al 2015; Danese et al 2009; Chapman et al. 2004.
- <sup>27</sup> Fuller-Thomson et al 2016; Dube et al. 2003.
- <sup>28</sup> Ford et al. 2011; Felitti et al. 1998.
- <sup>29</sup> Dube et al 2003; Dube et al. 2002.
- <sup>30</sup> Bryant et al. 2020; Dube et al 2003.
- <sup>31</sup> Choi et al. 2017.
- <sup>32</sup> Bledsoe et al. 2021; CDC 2019; Bethell et al. 2017; Hargreaves et al. 2017; Pachter et al. 2017; Verbitsky-Savitz 2016; Wade et al. 2016; Kagi and Regala 2012.
- <sup>33</sup> McRory et al. 2011; Shonkoff et al. 2009.
- <sup>34</sup> Bethell et al. 2017; Bledsoe et al. 2021; Centers for Disease Control and Prevention 2019; Hargreaves et al. 2017; Kagi and Regala 2012; Pachter et al. 2017; Shonkoff et al. 2009; Verbitsky-Savitz et al. 2016.

**Sociology 380 Researchers:** Destiny Love, Ke'ona Murrell, Katie Rose, and Melissa Slate  
**AREC Research Staff:** Mariah Benham, Erin Coulson, Holly Gavin, Elizabeth Gillenwater, Mitchell Henry, Jo Waiz  
**Primary Investigator:** Dr. Melissa S. Fry



# Adverse Childhood Experiences: Economy

2021 IU Southeast Sociology Research Lab

Adverse Childhood Experiences (ACEs) are traumatic events experienced from birth through age 17. Research shows that these experiences disrupt neurodevelopment, causing social, emotional, and cognitive impairments that affect behaviors, educational performance, and adult employment stability and income.<sup>1</sup>

Jefferson County has relatively low educational attainment, and the median household income is below the state median. The Jefferson County community-wide ACEs survey is an attempt to document the prevalence of ACEs and draw connections between experiences of childhood trauma and adult outcomes in the community.

## Key Findings

Based on a weighted sample:

- The estimated average ACE score for Jefferson County adults is 2.4.
- 30.4% of Jefferson County adults have an ACE score of 4 or higher.
- Roughly 9.7% of Jefferson County adults have an ACE score of 7 or higher.
- More than a third of Jefferson County adults report that before age 18 they experienced one or more of the following: separation or divorce

of parents, emotional abuse, or living with someone who suffered from substance use disorder (an alcoholic or drug user).

- 32.2% of Jefferson County adults have no ACEs.

## Employment and Income

- Those with higher income and educational attainment have lower ACE scores than those with lower income and educational attainment.<sup>2</sup>
- Among the mechanisms by which ACEs produce poorer outcomes in education and employment are the health and mental health outcomes that affect work performance and stability.<sup>3</sup>
- Research in Wales suggests the impact of ACEs on educational attainment is largely explained by failure to complete high school.<sup>4</sup>
- Research indicates that a combination of lower educational attainment and challenges with employment stability explain negative income outcomes for those with higher ACE scores.<sup>5</sup>

In the Jefferson County weighted sample, educational attainment is significantly associated with ACE scores.

Figure 1: Percent with Zero ACEs and Four or More ACEs that Report Having Struggled with Employment Stability (Sample N=637|Weighted N=24,391, p<.001)





Figure 2: Income Distribution for those with Zero ACEs and those With 4 or More ACEs, weighted sample (N=24,476, <001)



- Those with four or more ACEs are significantly less likely to have a college degree or higher, are less likely to have a high school diploma or equivalency, and are more likely to have less than high school (Figure 9).
- Among those who report 4 or more ACEs, 19.2% of the weighted sample reported that they had struggled to maintain consistent employment compared to 10.1% of the weighted sample with zero ACEs. The association is statistically significant (Figure 10,  $p < .001$ ).

Struggles with employment stability for those with higher ACE scores are likely the result of challenges with health, mental health, and substance abuse. When combined with lower educational attainment, these issues yield lower income levels. In the Jefferson County weighted sample the pattern is notable in higher percentages of those with four or more ACEs at lower income levels and lower percentages at the highest income levels than for those with zero ACEs. The significantly lower income levels of

those with higher ACE scores makes it more likely that children in their households will also experience chronic stress associated with poverty.

Effective and sustainable community and economic development efforts can reduce household financial stressors that contribute to the prevalence of ACEs. Prevention and mitigation priorities may include the following economic strategies<sup>34</sup>:

- Create and redesign policies and jobs that support workers with living wages.
- Build early care, school, and extracurricular youth programs that support families (birth to 18).

## References

<sup>1</sup>Felitti, Vincent J. et al. 1998; Anda, Robert F. et al. 2010. Verbitsky-Savitz et al. 2016.  
<sup>2</sup>Giano, Wheeler, and Hubach 2020.  
<sup>3</sup> DeVenter et al. 2020; Blodgett 2018; Hardcastle et al. 2018;  
<sup>4</sup> Hardcastle et al. 2018.  
<sup>5</sup>DeVenter et al. 2020; Metzler et al. 2017; Liu et al. 2012; Zielinski. 2009.  
<sup>34</sup> Bethell et al. 2017; Bledsoe et al. 2021; Centers for Disease Control and Prevention 2019; Hargreaves et al. 2017; Kagi and Regala 2012; Pachter et al. 2017; Shonkoff et al. 2009; Verbitsky-Savitz et al. 2016.



# Adverse Childhood Experiences: Education

2021 IU Southeast Sociology Research Lab

Adverse Childhood Experiences (ACEs) are traumatic events experienced from birth through age 17. Research shows that these experiences disrupt neurodevelopment, causing social, emotional, and cognitive impairments that affect behaviors, including health behaviors.<sup>1</sup> These impacts lead to negative outcomes in health, mental health, education, and social success and well being, ultimately resulting in early death.

## Key Findings

Based on a weighted sample:

- The estimated average ACE score for Jefferson County adults is 2.4.
- 30.4% of Jefferson County adults have an ACE score of 4 or higher.
- Roughly 9.7% of Jefferson County adults have an ACE score of 7 or higher.
- More than a third of Jefferson County adults report that before age 18 they experienced one or more of the following: separation or divorce of parents, emotional abuse, or living with someone who suffered from substance use disorder (an alcoholic or drug user).
- 32.2% of Jefferson County adults have no ACEs.

## ACES and Education


High ACE scores affect educational attainment.

Research indicates that high ACE scores are associated with the following:

- Below average literacy skills.
- Behavior problems that affect school performance.
- Frequent somatic symptoms when a teacher asks a child to do something.
- Departure from formal education with no qualifications.

In addition, behavior issues associated with ACEs can impact the classroom environment for all children.

Figure 1: Educational Attainment in Jefferson County, Indiana




Those with higher educational attainment have lower ACE scores than those with lower educational attainment.<sup>2</sup> Research suggests that among the mechanisms by which ACEs produce poorer outcomes in education and employment are the health and mental health outcomes that affect school engagement.<sup>3</sup>

Research in Wales finds the impact of ACEs on educational attainment is largely explained by failure to complete high school.<sup>4</sup>

Childhood trauma disrupts neurodevelopment, causing social, emotional, and cognitive impairments that affect behaviors.<sup>5</sup> These impacts lead to negative outcomes in several areas, including education.

Aggregated across a county, a higher than average prevalence of such experiences can result in lower than average educational attainment. Among Jefferson County residents age 25 and older, 16.3% have a bachelor's degree or higher compared with 32.7 percent nationally and 26.5% in the state of Indiana.

Figure 2: Educational Attainment for those with Zero ACEs and those with 4 or More ACEs, weighted sample ( $p < .001$ )



In the Jefferson County weighted sample, educational attainment is significantly associated with ACE scores. Those with four or more ACEs are significantly less likely to have a college degree or higher, are less likely to have a high school diploma or equivalency, and are more likely to have less than high school (Figure 2).

## Recommendations

While educational attainment is an outcome shaped by ACEs, it can also play an important role in prevention and mitigation of the impacts of ACEs. Priorities may include the following<sup>34</sup>:

- Create and redesign policies and jobs that support workers with living wages.
- Build early care, school, and extracurricular youth programs that support families (birth to 18).
- Teach social-emotional skills with attention to safe and healthy relationships and healthy parenting and family dynamics.
- Connect youth to caring adults.
- Provide widespread access to physical and

mental health care and family-centered treatment for substance use disorders.

The data from Jefferson County suggest significant opportunities to improve quality of life and community well-being through preventing and mitigating the impacts of childhood trauma. A focus on resilience and social support for healthy individuals and families can provide leverage for stronger and more sustainable community and economic development.


## References

- <sup>1</sup>Felitti, Vincent J. et al. 1998; Anda, Robert F. et al. 2010. Verbitsky-Savitz et al. 2016.
- <sup>2</sup>Giano, Wheeler, and Hubach 2020.
- <sup>3</sup>DeVenter et al. 2020; Blodgett 2018; Hardcastle et al. 2018;
- <sup>4</sup>Hardcastle et al. 2018.
- <sup>5</sup>McRory et al. 2011; Shonkoff et al. 2009.
- <sup>34</sup>Bethell et al. 2017; Bledsoe et al. 2021; Centers for Disease Control and Prevention 2019; Hargreaves et al. 2017; Kagi and Regala 2012; Pachter et al. 2017; Shonkoff et al. 2009; Verbitsky-Savitz et al. 2016.

Sociology 380 Researchers: Destiny Love, Ke'ona Murrell, Katie Rose, and Melissa Slate

AREC Research Staff: Mariah Benham, Erin Coulson, Holly Gavin, Elizabeth Gillenwater, Mitchell Henry, Jo Waiz





# Adverse Childhood Experiences: Health

2021 IU Southeast Sociology Research Lab

Adverse Childhood Experiences (ACEs) are traumatic events experienced from birth through age 17. Research shows that these experiences disrupt neurodevelopment, causing social, emotional, and cognitive impairments that affect behaviors, including health behaviors.<sup>1</sup> These impacts lead to negative outcomes in health, mental health, education, and social success and well being, ultimately resulting in early death.

Jefferson County is home to relatively high rates of smoking, obesity, physical inactivity, poor physical health days, and diabetes.

The Jefferson County community-wide ACEs survey documents the prevalence of ACEs and draws important connections between experiences of childhood trauma and health outcomes.

## Key Findings

Based on a weighted sample:

- The estimated average ACE score for Jefferson County adults is 2.4.
- 30.4% of Jefferson County adults have an ACE score of 4 or higher.
- Roughly 9.7% of Jefferson County adults have an ACE score of 7 or higher.
- More than a third of Jefferson County adults report that before age 18 they experienced one

or more of the following: separation or divorce of parents, emotional abuse, or living with someone who suffered from substance use disorder (an alcoholic or drug user).

- 32.2% of Jefferson County adults have no ACEs.

## Health Behaviors and Outcomes

Those with higher ACE scores experience progressively higher rates of disease and dysfunction in their adult lives. Those with zero reported ACEs have significantly better outcomes than those with four or more ACEs across a range of indicators of adult well-being and predictors of early death.<sup>3</sup>

- Toxic stress is one mechanism by which childhood trauma affects adult health outcomes.
- The body experiences chronic stressors and remains in a fight or flight mode of vigilance that causes the release of hormones and chemicals that affect neurodevelopment and long-term health outcomes.<sup>12</sup>
- Chronic stress causes changes to neurobiology that impact early brain development,<sup>13</sup> the immune system,<sup>14</sup> and the endocrine system.<sup>15</sup>
- Stressful or traumatic experiences often lead to

Figure 12: Percent Indicating they Smoke or Have Ever Regularly Smoked Cigarettes by ACE score category (Sample N=642 | Weighted N=24,474, p<.001)





Figure 13: Prevalence of Health Issues for those with Zero ACEs and those with Four or More ACEs in Jefferson County, IN (Sample N=438 | Weighted N=25,567)



Statistically significant differences in prevalence between those with zero ACE and those with four or more ACE are indicated with asterisks. \*p<.05, \*\*p<.01, \*\*\*p<.001. Contrary to findings in the literature, associations with liver disease and obesity were not found to be statistically significant in the Jefferson County sample, with or without weighting.

social, emotional, and cognitive deficiencies that increase risk for unhealthy behaviors and chronic disease.<sup>16</sup>

- Toxic stress can cause issues with immune and metabolic systems that fight illness, leading to a lifetime of susceptibility to illness.<sup>17</sup>
- Toxic stress causes early onset of disease, disability, and premature death.<sup>18</sup>
- ACE exposure increases risk of obesity.<sup>19</sup>
- Higher ACE scores are associated with increased risk of smoking, heart disease, chronic lung disease, and poor self-rated health.
- Those with 4 or more ACEs are more likely to report high numbers of sexual partners and incidence of sexually transmitted disease (2-4 fold increase in risks).<sup>20</sup>
- Researchers find a 1.4-1.6 fold increase in physical inactivity and severe obesity among those with 4 or more ACEs compared to those with none.<sup>21</sup>
- Higher ACE scores are associated with diseases such as ischemic heart disease, cancer, chronic lung disease, skeletal fractures, and liver disease.<sup>22</sup> Research links ACEs to systemic arterial stiffness among adolescents.<sup>23</sup>

## Jefferson County

Jefferson County residents report an average 4.2 poor physical health days per month compared to 4.0 for the state and 3.4 among U.S. top performers. The county has relatively high rates of smoking (24%), obesity (33%), and physical inactivity (25%). In Jefferson County,

- Those with four or more ACEs are significantly more likely than those with zero ACEs to smoke or to have smoked regularly at some point in their life (Figure 1).
- A higher ACE score is significantly associated with higher rates of sexually transmitted infections, heart disease, chronic lung disease, and physical disability that limits movement (Figure 2).
- Obesity rates were not significantly different between those with zero and those with four or more ACEs.

Findings from Jefferson County largely confirm the existing research on the association between ACEs and adult health outcomes. With a higher average ACE score than the nation, it is not surprising that Jefferson County also sees higher rates of chronic disease.

\*Endnote numbers match the full report available at [www.resilientjeffersoncounty.org](http://www.resilientjeffersoncounty.org). Full citations also available on the website.



# Adverse Childhood Experiences: Substance Abuse and Mental Health

2021 IU Southeast Sociology Research Lab

Adverse Childhood Experiences (ACEs) are traumatic events experienced from birth through age 17. Research shows that these experiences disrupt neurodevelopment, causing social, emotional, and cognitive impairments that affect behaviors, including health behaviors.<sup>1</sup> These impacts lead to negative outcomes in health, mental health, education, and social success and well being, ultimately resulting in early death.

Jefferson County is home to relatively high rates of poor mental health days and far higher suicide rate than the state or nation.

The Jefferson County community-wide ACEs survey is an attempt to document the prevalence of ACEs, draw connections between experiences of childhood trauma and health, mental health, education, and economic outcomes, and identify the highest priorities for prevention and intervention. This fact sheet focuses on connections to substance abuse and mental health.

## Key Findings on ACEs

Based on a weighted sample:

- The estimated average ACE score for Jefferson

County adults is 2.4.

- 30.4% of Jefferson County adults have an ACE score of 4 or higher.
- Roughly 9.7% of Jefferson County adults have an ACE score of 7 or higher.
- More than a third of Jefferson County adults report that before age 18 they experienced one or more of the following: separation or divorce of parents, emotional abuse, or living with someone who suffered from substance use disorder (an alcoholic or drug user).
- 32.2% of Jefferson County adults have no ACEs.

## ACEs, Mental Health, & Substance Use Disorder

Childhood trauma increases risk for a range of mental health concerns and for substance use disorder (SUD). Bryant (2020) found that each individual ACE type was significantly associated with SUD.<sup>2</sup> These issues may impact employment stability. Studies find associations between early adversity and each of the following:

Figure 1: Prevalence of Reported Mental Health Issues for those with Zero ACEs Compared to those with 4 or More ACEs





Figure 2: Percent Indicating any Substance Use Disorder and Misuse of Prescription Painkillers or Use of Street Opioids Such as Heroin among those with Zero ACEs and those with Four or More ACEs



- Frequent mental distress.<sup>3</sup>
- Depression.<sup>4</sup>
- Suicide attempts.<sup>5</sup>
- Smoking.<sup>6</sup>
- Alcohol abuse.<sup>7</sup>
- Substance abuse.<sup>8</sup>

In addition, sexual abuse and parental/other family member’s mental illness increase the odds for having a suicide attempt for both men and women and emotional neglect is also a factor for men.<sup>9</sup>

## ACEs, Mental Health & Substance Abuse in Jefferson County, IN

County Health Rankings data indicate that Jefferson County residents report an average 5.2 poor mental health days per month compared to 4.7 for the state and 3.8 among U.S. top performers.

In Jefferson County, as one moves up in ACE score category, so too does the likelihood of depression, anxiety, and suicidal thoughts. The ACEs survey revealed that those who report four or more ACEs are significantly more likely than those with zero ACEs to report:

- A history of depression.
- Struggling with anxiety.
- Experiencing suicidal thoughts.

Finally, in a state and region beset with some of the worst of the opioid epidemic, when

compared to those with zero ACEs, those with four or more ACEs are

- 10 times more likely to report a diagnosis or self-assessment of substance use disorder.
- To have ever misused prescription pain killers or to have used street opioids such as heroin (Figure 16).

Findings from Jefferson County confirm the existing research on the association between ACEs and adult mental health and substance abuse outcomes. With a higher average ACE score than the nation, it is not surprising that Jefferson County also sees higher rates of mental health challenges and substance use disorder.

## References

- <sup>1</sup>Felitti, Vincent J. et al. 1998; Anda, Robert F. et al. 2010. Verbitsky-Savitz et al. 2016.
- <sup>2</sup>Bryant, Coman, and Damian 2020.
- <sup>3</sup>Gilbert et al 2015.
- <sup>4</sup>Gilbert et al 2015; Danese et al 2009; Chapman et al. 2004.
- <sup>5</sup>Fuller-Thomson et al 2016; Dube et al. 2003.
- <sup>6</sup>Ford et al. 2011; Felitti et al. 1998.
- <sup>7</sup>Dube et al 2003; Dube et al. 2002.
- <sup>8</sup>Bryant et al. 2020; Dube et al 2003.
- <sup>9</sup>Choi et al. 2017.

Sociology 380 Researchers: Destiny Love, Ke’ona Murrell, Katie Rose, and Melissa Slate  
 AREC Research Staff: Mariah Benham, Erin Coulson, Holly Gavin, Elizabeth Gillenwater, Mitchell Henry, Jo Waiz  
 Primary Investigator: Dr. Melissa S. Fry